

Crystal Data: Orthorhombic. *Point Group:* $2/m\ 2/m\ 2/m$. As rims to 10 μm on holtite, in the outer portion of the cores of zoned As- and Sb-dumortierite crystals.

Physical Properties: *Cleavage:* n.d. *Fracture:* n.d. *Tenacity:* n.d. *Hardness:* = n.d. $D(\text{meas.}) = \text{n.d.}$ $D(\text{calc.}) = 3.72$

Optical Properties: Translucent. *Color:* Creamy white to brownish yellow or gray-yellow. *Streak:* White. *Luster:* n.d. *Optical Class:* Biaxial (-). $\alpha = 1.740\text{--}1.747$ β and $\gamma = \sim 1.76$

Cell Data: *Space Group:* $Pnma$. [By analogy to dumortierite.] $a = \sim 4.7001$ $b = \sim 11.828$ $c = \sim 20.243$ $Z = 4$

X-ray Powder Pattern: Calculated pattern.

3.2305 (100), 10.21 (67), 5.8610 (66), 2.9305 (65), 2.8945 (64), 3.4582 (63), 3.0675 (53)

Chemistry:	(1)
P_2O_5	0.26
Nb_2O_5	5.21
Ta_2O_5	0.66
SiO_2	18.68
TiO_2	0.11
B_2O_3	4.91
Al_2O_3	49.74
As_2O_3	5.92
Sb_2O_3	10.81
FeO	0.51
H_2O	[0.05]
Total	96.86

(1) Marta mine, Szklana Hill, Lower Silesia, Poland; average electron microprobe analysis, H_2O calculated for charge balance; corresponds to $\{(\text{Nb}_{0.26}\text{Ta}_{0.02}\square_{0.18})(\text{Al}_{0.27}\text{Fe}_{0.05}\text{Ti}_{0.01})\square_{0.21}\}_{\Sigma=1.00}$ $\text{Al}_6\text{B}_{0.92}\{\text{Si}_{2.03}\text{P}_{0.02}(\text{Sb}_{0.48}\text{As}_{0.39}\text{Al}_{0.07})\}_{\Sigma=3.00}(\text{O}_{17.09}\text{OH}_{0.04}\square_{0.87})_{\Sigma=18.00}$.

Mineral Group: Holtite group, dumortierite supergroup.

Occurrence: In the internal portion of a complex zoned granitic pegmatite.

Association: Holtite, microcline, quartz, muscovite, spessartine, chrysoberyl, zircon, monazite-(Ce), cheralite, xenotime-(Y), Mn-rich fluor-, hydroxyl- and chlorapatite, beusite, columbite-(Fe), columbite-(Mn), tantalite-(Mn), stibiocolumbite, stibiotantalite, fersmite, pyrochlore-supergroup minerals, and other minerals.

Distribution: From the Marta mine, northern part of Szklana Hill, Szklary serpentinite massif, ~60 km south of Wrocław, Lower Silesia, Poland.

Name: For the composition and the relationship to *holtite*.

Type Material: Mineralogical Museum, University of Wrocław, Faculty of Earth Science and Environmental Management, Institute of Geological Sciences, Poland (MMWr IV7615). Also at the National Museum of Natural History (Smithsonian Institution), Washington, D.C., USA (NMNH 175986-175988).

References: (1) Pieczka, A., R.J. Evans, E.S. Grew, L.A. Groat, C. Ma, and G.R. Rossman (2013) The dumortierite supergroup. II. Three new minerals from the Szklary pegmatite, SW Poland: Nioboholtite, $(\text{Nb}_{0.6}\square_{0.4})\text{Al}_6\text{BSi}_3\text{O}_{18}$, titanoholtite, $(\text{Ti}_{0.75}\square_{0.25})\text{Al}_6\text{BSi}_3\text{O}_{18}$, and szklaryite, $\square\text{Al}_6\text{BAAs}^{3+}_3\text{O}_{15}$. *Mineral. Mag.*, 77(6), 2841-2856. (2) (2015) *Amer. Mineral.*, 100, 2012-2013 (abs. ref. 1).